arXiv:1510.00318v2 [math.RA] 30 Mar 2016

About special elements in quaternion algebras over finite fields

Diana SAVIN

Abstract. In this paper we study special Fibonacci quaternions and special generalized Fibonacci-Lucas quaternions in quaternion algebras over finite fields.

Key Words: quaternion algebras; finite fields; Fibonacci numbers; Lucas numbers, Fibonacci quaternions, generalized Fibonacci-Lucas quaternions.

2000 AMS Subject Classification: 15A06, 16G30, 1R52, 11B39.

1. Introduction

Let K be an algebraic number field and let $\mathbb{H}_K(\alpha,\beta)$ *be the generalized quaternion algebra* over the field K. An arbitrary element from this algebra has the form $x = x_1 \cdot 1 + x_2 e_2 + x_3 e_3 + x_4 e_4$, where $x_i \in K, i \in \{1,2,3,4\}$, where the elements of the basis $\{1, e_2, e_3, e_4\}$ satisfy the following multiplication table:

$\cdot$	1	e_2	e_3	e_4
1	1	e_2	e_3	e_4
e_2	e_2	α	e_4	αe_3
e_3	e_3	$-e_4$	β	$-\beta e_2$
e_4	e_4	$-\alpha e_3$	βe_2	$-\alpha\beta$

Let $x \in \mathbb{H}_K(\alpha,\beta)$ and let $\boldsymbol{n}(x)$ be the norm of a generalized quaternion x. This norm has the following expression $\boldsymbol{n}(x) = x_1^2 - \alpha x_2^2 - \beta x_3^2 + \alpha\beta x_4^2$.We recall that $\mathbb{H}_K(\alpha,\beta)$ is a division algebra if and only if for $x \in \mathbb{H}_K(\alpha,\beta)$ we have $\boldsymbol{n}(x) = 0$ if and only if $x = 0$. Otherwise, the algebra $\mathbb{H}_K(\alpha,\beta)$ is called *a split* algebra.
When $\alpha = \beta = -1$ and $K = \mathbb{R}$ we obtain Hamilton quaternion algebra $\mathbb{H}_{\mathbb{R}}(-1,-1)$, with the basis $\{1,i,j,k\}$. It is known that this algebra is a division algebra.
When $\alpha = \beta = -1$ and K is a finite field ($K = \mathbb{Z}_p$, where p is a prime positive integer) we obtain the quaternion algebra $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$. It is known that this algebra is a split algebra ([Gr, Mi, Ma; 15], Theorem 2, p. 882).

Let $(f_n)_{n\geq 0}$ be the Fibonacci sequence:

$$f_0 = 0; f_1 = 1; f_n = f_{n-1} + f_{n-2},\ n \geq 2$$

and $(l_n)_{n\geq 0}$ be the Lucas sequence:

$$l_0 = 2; l_1 = 1; l_n = l_{n-1} + l_{n-2},\ n \geq 2$$

In his paper [Ho; 63], A. F. Horadam introduced the Fibonacci quaternions $(F_n)_{n\geq 0}$:

$$F_n = f_n + f_{n+1} \cdot i + f_{n+2} \cdot j + f_{n+3} \cdot k,\ n \geq 0.$$

Some properties of Fibonacci quaternions in certain quaternion algebras were investigated in the papers [Ak, Ko, To; 14], [Ta; Ya; 15], [Po, Ki, Ke; 16], [Yü, Ay; 16], etc. In this paper we determine the Fibonacci quaternions (respectively generalized Fibonacci- Lucas quaternions) which are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$ and the Fibonacci quaternions which are invertible in the quaternion ring $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$.
We recall some necessary results about Fibonacci sequence and about Fibonacci quaternions.

Proposition 1.1 ([Fib]). *Let $(f_n)_{n\geq 0}$ be the Fibonacci sequence. If f_n is a prime number ($n \geq 5$), then n is a prime number.*

Proposition 1.2 ([Fib]). *Let $(f_n)_{n\geq 0}$ be the Fibonacci sequence. If $n, m \in \mathbb{N}^*$, $n|m$, then $f_n|f_m$.*

Theorem 1.1 ([Su], [Su; 92]). *Let $p \notin \{2,5\}$ be a prime positive integer. Then*

$$f_{\frac{p-\left(\frac{p}{5}\right)}{2}} \equiv \begin{cases} 0\,(mod\,p), if\ p \equiv 1(mod 4), \\ 2\,(-1)^{\left[\frac{p+5}{10}\right]} \cdot \left(\frac{p}{5}\right) \cdot 5^{\frac{p-3}{4}}\,(mod\,p), \text{ if } p \equiv 3(mod 4) \end{cases}$$

where $[x]$ is the integer part of x and $\left(\frac{p}{5}\right)$ is the Legendre symbol.

Proposition 1.3 ([Fib]). *The cycle of the Fibonacci numbers mod 2 is*

$$0, 1, 1(0, 1, 1), ...$$

so:
i) *the cycle-length of the Fibonacci numbers mod 2 is 3;*
ii) *a Fibonacci number f_n is even if and only if $n \equiv 0\ (mod\ 3)$.*
iii) *a Lucas number l_n is even if and only if $n \equiv 0\ (mod\ 3)$.*

Proposition 1.4 ([Fib]). *The cycle of the Fibonacci numbers mod 3 is*

$$0, 1, 1, 2, 0, 2, 2, 1(0, 1, 1, 2, 0, 2, 2, 1), ...$$

so, the cycle-length of the Fibonacci numbers mod 3 *is* 8.

Proposition 1.5 ([Fib]). *The cycle of the Fibonacci numbers mod* 5 *is*

$$0,1,1,2,3,0,3,3,1,4,0,4,4,3,2,0,2,2,4,1(0,1,1,2,3,0,3,3,1,4,0,4,4,3,2,0,2,2,4,1),...$$

so:
i) *the cycle-length of the Fibonacci numbers mod* 5 *is* 20;
ii) $f_n \equiv 0 \ (mod\ 5)$ *if and only if* $n \equiv 0 \ (mod\ 5)$.

Proposition 1.6 ([Ho; 63]). *Let* $(f_n)_{n\geq 0}$ *be the Fibonacci sequence and* $F_n (n \geq 0)$ *be the Fibonacci quaternions. Then, the norm of the nth Fibonacci quaternions is:*

$$n(F_n) = 3f_{2n+3}.$$

Let p be an odd prime positive integer. In the paper [Mi, Se; 11], C. J. Miguel, R. Serodio calculated the number of zero-divisors in $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$. They proved that:

Theorem 1.2 ([Mi, Se; 11]). *The number of zero-divisors in* $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$ *is*:

$$p^3 + p^2 - p.$$

Let p be a prime positive integer. According to [Da, Dr; 70], if f_z is the smallest non-zero Fibonacci number with the property $p|f_z$, then $z = z(p)$ is defined as the entry point of p in the Fibonacci sequence.

Remark 1.1. ([Da, Dr; 70]). $p|f_n$ *if and only if* $z(p)|n$, $(\forall)\, n \in \mathbb{N}$.

Definition 1.1. ([Da, Dr; 70]). *Let* $(f_n)_{n\geq 0}$ *be the Fibonacci sequence. If we reduce the Fibonacci sequence modulo* p, *to obtain a periodic sequence. The period* $k = k(p)$ *is the smallest integer* k *for which*

$$f_k \equiv 0 \ (mod\ p) \ \text{ and } \ f_{k+1} \equiv 1 \ (mod\ p).$$

For example: $k(2) = z(2) = 3$, $k(3) = 8$, $z(3) = 4$, $k(5) = 20$, $z(5) = 5$ (according to Proposition 1.3, Proposition 1.4, Proposition 1.5).

Remark 1.2. ([Da, Dr; 70]). *The entry point* $z(p)$ *divides the period* $k(p)$.

Remark 1.3. ([Da, Dr; 70]). *Let* p *be an odd prime positive integer. The following statements are true:*

i) $k(p) = z(p)$ *if* $z(p) \equiv 2 \ (mod\ 4)$;
ii) $k(p) = 2z(p)$ *if* $z(p) \equiv 0 \ (mod\ 4)$;
iii) i) $k(p) = 4z(p)$ *if* $z(p)$ *is odd.*

Proposition 1.7. ([Fib]).*Let* $(f_n)_{n\geq 0}$ *be the Fibonacci sequence* and $(l_n)_{n\geq 0}$ *be the Lucas sequence Then:*
i)
$$f_n^2 + f_{n+1}^2 = f_{2n+1}, \forall\ n \in \mathbb{N};$$
ii)
$$f_n + f_{n+4} = 3f_{n+2}, \forall\ n \in \mathbb{N};$$
iii)
$$l_n^2 + l_{n+1}^2 = 5f_{2n+1}, \forall\ n \in \mathbb{N};$$
iv)
$$f_m l_{m+p} = f_{2m+p} + (-1)^{m+1} \cdot f_p, \forall\ m, p \in \mathbb{N};$$
v)
$$f_n + f_{n+2} = l_{n+1}, \forall\ n \in \mathbb{N};$$

2. Special Fibonacci quaternions in the quaternion algebra $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$.

In this section we determine the zero divisors Fibonacci quaternions, respectively the invertible Fibonacci quaternions in the quaternion rings $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$, $\mathbb{H}_{\mathbb{Z}_3}(-1,-1)$, $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$, $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$, where p is a prime positive integer, $p > 5$. The idea for to find these elements is the following: since $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$, (where p is a prime positive integer) is a split algebra, it is isomorphic to the matrix ring $M_2(\mathbb{Z}_p)$. If $\varphi : \mathbb{H}_{\mathbb{Z}_p}(-1,-1) \to M_2(\mathbb{Z}_p)$ is a rings isomorphism, then a Fibonacci quaternion F_n is a zero divisor in $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$ if and only if $\varphi(F_n)$ is a zero divisor in $M_2(\mathbb{Z}_p)$ (see [Mi, Se; 11], Theorem 2.1). Moreover, it is know that a matrix $A \in M_2(\mathbb{Z}_p)$ is a zero divisor in $M_2(\mathbb{Z}_p)$ if and only if it is not invertible in $M_2(\mathbb{Z}_p)$. These involve that, a Fibonacci quaternion is a zero divisor in $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$ if and only if it is not invertible in $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$. So, a Fibonacci quaternion F_n is a zero divisor in $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$ if and only if $n(F_n) = 0$.

Proposition 2.1. i) *Fibonacci quaternions* F_n *are zero divisors in the quaternion ring* $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$ *if and only if* $n \equiv 0 \ (mod\ 3)$.
ii) *In* $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$ *there are one Fibonacci quaternion which is zero divisor and two Fibonacci quaternions which are invertible.*

Proof. i) According to Proposition 1.6 and Proposition 1.3 we obtain that Fibonacci quaternions F_n are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$ if

and only if $f_{2n+3} \equiv 0 \pmod 2$ if and only if $n \equiv 0 \pmod 3$.
ii) There are 16 elements in the quaternion algebra $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$. According to Theorem 1.2 we know that there are 10 zero divisors in $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$. We want to determine how many of them are zero divisors Fibonacci quaternions and after this we determine how many invertible Fibonacci quaternions are in the quaternion algebra $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$.
Applying i) and Proposition 1.3 we obtain that there is one Fibonacci quaternion which is zero divisor in $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$, namely for $f_n = \overline{0}$, $f_{n+1} = \overline{1}$, $f_{n+2} = \overline{1}$, $f_{n+3} = \overline{0}$ (in $\mathbb{Z}_2$), so $F_n = i + j$.
Applying i) and Proposition 1.3, it results immediately that there are two invertible Fibonacci quaternions in $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$: $F_{n_1} = \overline{1} + i + k$ and $F_{n_2} = \overline{1} + j + k$.

Proposition 2.2. *All Fibonacci quaternions F_n are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_3}(-1,-1)$ and the number of these elements is* 8.

Proof. Applying Proposition 1.6 we obtain that $n(F_n) = \overline{0}$ in $\mathbb{Z}_3$, $(\forall)\, n \in \mathbb{N}$. This implies that all Fibonacci quaternions F_n are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_3}(-1,-1)$.
There are 81 elements in the quaternion algebra $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$. According to Theorem 1.2, 33 of them are zero divisors. Now, we determine how many of these 33 elements are zero divisors Fibonacci quaternions. From the previously proved and Proposition 1.4 we obtain the following zero divisors Fibonacci quaternions in $\mathbb{H}_{\mathbb{Z}_3}(-1,-1)$:
$F_{n_1} = i + j + \overline{2}k$, $F_{n_2} = \overline{1} + i + \overline{2}j$, $F_{n_3} = \overline{1} + \overline{2}i + \overline{2}k$, $F_{n_4} = \overline{2} + \overline{2}j + \overline{2}k$, $F_{n_5} = \overline{2}i + \overline{2}j + k$, $F_{n_6} = \overline{2} + \overline{2}i + j$, $F_{n_7} = \overline{2} + i + k$, $F_{n_8} = \overline{1} + j + k$.

Proposition 2.3. i) *Fibonacci quaternions F_n are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$ if and only if $n \equiv 1$ (mod* 5).
ii) *In $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$ there are* 4 *Fibonacci quaternions which are zero divisors and* 16 *Fibonacci quaternions which are invertible.*

Proof. i) According to Proposition 1.6 and Proposition 1.5 (ii) we obtain that Fibonacci quaternions F_n are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$ if and only if $f_{2n+3} \equiv 0$ (mod 5) if and only if $2n + 3 \equiv 0 \pmod 5$ if and only if $n \equiv 1 \pmod 5$.
ii) There are 625 elements in the quaternion algebra $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$. According to Theorem 1.2, 145 of them are zero divisors. Applying i) and Proposition 1.5 we obtain 4 zero divisor Fibonacci quaternion in $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$:
$F_{n_1} = \overline{1} + i + \overline{2}j + \overline{3}k$, $F_{n_2} = \overline{3} + \overline{3}i + j + \overline{4}k$, $F_{n_3} = \overline{4} + \overline{4}i + \overline{3}j + \overline{2}k$, $F_{n_4} = \overline{2} + \overline{2}i + \overline{4}j + k$.
Applying i) and Proposition 1.5, it results immediately that there are 16 invertible Fibonacci quaternions in $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$: $F_{n_1} = i + j + \overline{2}k$, $F_{n_2} = \overline{1} + \overline{2}i + \overline{3}j$, $F_{n_3} = \overline{2} + \overline{3}i + \overline{3}k$, $F_{n_4} = \overline{3} + \overline{3}j + \overline{3}k$, $F_{n_5} = \overline{3}i + \overline{3}j + k$, $F_{n_6} = \overline{3} + i + \overline{4}j$, $F_{n_7} = \overline{1} + \overline{4}i + \overline{4}k$, $F_{n_8} = \overline{4} + \overline{4}j + \overline{4}k$, $F_{n_9} = \overline{4}i + \overline{4}j + \overline{3}k$, $F_{n_{10}} = \overline{4} + \overline{3}i + \overline{2}j$, $F_{n_{11}} = \overline{3} + \overline{2}i + \overline{2}k$, $F_{n_{12}} = \overline{2} + \overline{2}j + \overline{2}k$, $F_{n_{13}} = \overline{2}i + \overline{2}j + \overline{4}k$, $F_{n_{14}} = \overline{2} + \overline{4}i + j$, $F_{n_{15}} = \overline{4} + i + k$, $F_{n_{16}} = \overline{1} + j + k$.

We asked our self what happens in the general case, when we work in the quaternion ring $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$. We obtain the following result:

Theorem 2.1. *Let p be a prime positive integer, $p\equiv 13,17 \pmod{20}$, l be a positive integer and $n=\frac{p\cdot(2l+1)+2l-1}{4}-1$. Then, the Fibonacci quaternions F_n are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$.*

Proof. We remark that $4|(p\cdot(2l+1)+2l-1)$, so n is an integer number. Since $p\equiv 13,17 \pmod{20}$, it results that the Legendre symbol $\left(\frac{p}{5}\right)=-1$ and $\frac{p-\left(\frac{p}{5}\right)}{2}=\frac{p+1}{2}$ is an odd integer number. Since $n=\frac{p\cdot(2l+1)+2l-1}{4}-1$, it results $\frac{p+1}{2}|(2n+3)$. Applying Proposition 1.2, it results that $f_{\frac{p+1}{2}}|f_{2n+3}$. According to Theorem 1.1 and Proposition 1.6, we obtain that $n(F_n)=\overline{0}$ in $\mathbb{Z}_p$. So, if p is a prime positive integer, $p\equiv 13,17 \pmod{20}$ and $n=\frac{p\cdot(2l+1)+2l-1}{4}-1$, $l\in\mathbb{N}^*$, then the Fibonacci quaternions F_n are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$.

Now, we generalize the results from Proposition 2.1 and Proposition 2.3, obtaining a necessary and sufficient condition for that a Fibonacci quaternion F_n to be a zero divisor in the quaternion ring $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$. The difference between the expression of n from Theorem 2.1 and the expression of n from Theorem 2.2 is that n from Theorem 2.2 depends of $z(p)$.

Theorem 2.2. *Let p be a prime positive integer, $p>5$. Then*:
i) *Fibonacci quaternions F_n are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$ if and only if $n=\frac{(2l+1)\cdot z(p)-1}{2}-1$, where $l\in\mathbb{N}$.*
ii) *In the quaternion ring $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$ there are 4 Fibonacci quaternions which are zero divisors and $k(p)-4$ Fibonacci quaternions which are invertible.*

Proof. i) Let F_n be a Fibonacci quaternion which is zero divisor in the quaternion ring $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$. Applying Proposition 1.6, it results that $p|f_{2n+3}$. This is equivalent with $z(p)\,|\,(2n+3)$ (according to Remark 1.1). This is equivalent with $z(p)$ is an odd number and $(\exists)\ l\in\mathbb{N}$ such that $2n+3=(2l+1)\cdot z(p)$. We obtain that $n=\frac{(2l+1)\cdot z(p)-1}{2}-1$, $l\in\mathbb{N}$.
ii) Since $z(p)$ is an odd number, using Remark 1.3 and a reasoning similar to the one used in Proposition 2.1 (ii) and Proposition 2.3 (ii), we obtain that Fibonacci quaternions F_n are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$ if and only if $f_{2n+3}\equiv 0 \pmod{p}$ if and only if $2n+3\equiv 0 \pmod{z(p)}$(according to Remark 1.1) if and only if $2n\equiv z(p)-3 \pmod{z(p)}$. Since $g.c.d.(2,z(p))=1$, it results that $\overline{2}$ is invertible in $\mathbb{Z}_{z(p)}$, so, the congruence $2n\equiv z(p)-3 \pmod{z(p)}$ has unique solution mod $z(p)$. Applying Remark 1.3, we get that in $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$ there are 4 Fibonacci quaternions which are zero divisors and $k(p)-4$ Fibonacci quaternions which are invertible.

3. Generalized Fibonacci-Lucas quaternions which are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_r}(-1,-1)$.

Let $(f_n)_{n\geq 0}$ be the Fibonacci sequence and $(l_n)_{n\geq 0}$ be the Lucas sequence.

In the paper [Fl, Sa; 15], we introduced the generalized Fibonacci-Lucas numbers and the generalized Fibonacci-Lucas quaternions $(F_n)_{n\geq 0}$, namely: let n,p,q be arbitrary integers, $n\geq 0$. The numbers $(g_n^{p,q})_{n\geq 0}$,

$$g_{n+1}^{p,q}=pf_n+ql_{n+1},\ n\geq 0$$

are called *the generalized Fibonacci-Lucas numbers.*

Let r be a prime positive integer and let $\mathbb{H}_{\mathbb{Z}_r}(-1,-1)$ be the quaternion algebra over the finite field $\mathbb{Z}_r$ with a basis $\{1,i,j,k\}$, where $i^2=-1$, $j^2=-1$, $k=ij=-ji$. The element

$$G_n^{p,q}=g_n^{p,q}+g_{n+1}^{p,q}\cdot i+g_{n+2}^{p,q}\cdot j+g_{n+3}^{p,q}\cdot k$$

is called *the n-th generalized Fibonacci-Lucas quaternion* (in the quaternion algebra $\mathbb{H}_{\mathbb{Z}_r}(-1,-1)$).

We determine the generalized Fibonacci-Lucas quaternions which are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_r}(-1,-1)$, when $r=2$ or $r=3$ or $r=5$ or $r=p$ or $r=q$.

First, we compute the norm for the n-th generalized Fibonacci-Lucas quaternion in $\mathbb{H}_{\mathbb{Z}_r}(-1,-1)$.

Proposition 3.1. *Let n,p,q be arbitrary integers, $n\geq 0$. Let $G_n^{p,q}$ be the n-th generalized Fibonacci-Lucas quaternion. Let r be a prime positive integer. Then, the norm of $G_n^{p,q}$ in the quaternion algebra $\mathbb{H}_{\mathbb{Z}_r}(-1,-1)$, is given by*

$$\mathbf{n}(G_n^{p,q})=\overline{3p^2f_{2n+1}+15q^2f_{2n+3}+6pq(f_{2n+1}+f_{2n+3})}\ (in\ \mathbb{Z}_r).$$

Proof.

$$n(G_n^{p,q})=G_n^{p,q}\overline{G_n^{p,q}}=(g_n^{p,q})^2+\left(g_{n+1}^{p,q}\right)^2+\left(g_{n+2}^{p,q}\right)^2+\left(g_{n+3}^{p,q}\right)^2=$$

$$=(pf_{n-1}+ql_n)^2+(pf_n+ql_{n+1})^2+(pf_{n+1}+ql_{n+2})^2+(pf_{n+2}+ql_{n+3})^2=$$
$$=p^2\left(f_{n-1}^2+f_n^2+f_{n+1}^2+f_{n+2}^2\right)+q^2\left(l_n^2+l_{n+1}^2+l_{n+2}^2+l_{n+3}^2\right)+$$
$$+2pq\left(f_{n-1}l_n+f_nl_{n+1}+f_{n+1}l_{n+2}+f_{n+2}l_{n+3}\right).$$

Applying Proposition 1.7 (i, iii, iv), we have:

$$n(G_n^{p,q})=p^2(f_{2n-1}+f_{2n+3})+q^2(5f_{2n+1}+5f_{2n+5})+$$

$$+2pq\left[f_{2n-1}+(-1)^n\cdot f_1+f_{2n+1}+(-1)^{n+1}\cdot f_1+f_{2n+3}+(-1)^{n+2}\cdot f_1+f_{2n+5}+(-1)^{n+3}\cdot f_1\right].$$

Using Proposition 1.7 (ii), we obtain:

$$\mathbf{n}\left(G_n^{p,q}\right) = \overline{3p^2 f_{2n+1} + 15q^2 f_{2n+3} + 6pq\left(f_{2n+1} + f_{2n+3}\right)} \ (in\ \mathbb{Z}_r).$$

Proposition 3.2. *Let p, q be two integers and let n be an arbitrary positive integer. Then, the following statements are true:*
i) *if $p \equiv 0\ (mod\ 2)$ and $q \equiv 1\ (mod\ 2)$, then the generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$ if and only if $n \equiv 0\ (mod\ 3)$;*
ii) *if $p \equiv 1\ (mod\ 2)$ and $q \equiv 0\ (mod\ 2)$ then the generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$ if and only if $n \equiv 1\ (mod\ 3)$;*
iii) *if $p; q \equiv 0\ (mod\ 2)$ then all generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$;*
iv) *if $p; q \equiv 1\ (mod\ 2)$, then the generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$ if and only if $n \equiv 2\ (mod\ 3)$;*

Proof. i) The generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$ if and only if $n\left(G_n^{p,q}\right) = \overline{0}$ (in $\mathbb{Z}_2$). Applying Proposition 3.1 and the facts that $r = 2$ and $p \equiv 0 \pmod 2$ and $q \equiv 1 \pmod 2$, it results that $n\left(G_n^{p,q}\right) = \overline{f_{2n+3}}$ in $\mathbb{Z}_2$. Applying Proposition 1.3, $\overline{f_{2n+3}} = \overline{0}$ in $\mathbb{Z}_2$ if and only if $n \equiv 0 \pmod 3$.
ii) If $p \equiv 1 \pmod 2$ and $q \equiv 0 \pmod 2$, similarly as in i) we get that the generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$ if and only if $\overline{f_{2n+1}} = \overline{0}$ in $\mathbb{Z}_2$ if and only if $n \equiv 1 \pmod 3$.
iii) If $p; q \equiv 0 \pmod 2$, it results that $n\left(G_n^{p,q}\right) = \overline{0}$ (in $\mathbb{Z}_2$) for $(\forall)\ G_n^{p,q}$, so, we obtain immediate that all generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$.
iv) If $p; q \equiv 1 \pmod 2$, applying Proposition 1.7 (v) it results that $n\left(G_n^{p,q}\right) = \overline{f_{2n+1}} + \overline{f_{2n+3}} = \overline{l_{2n+2}}$ (in $\mathbb{Z}_2$). Using a reasoning as in i) and Proposition 1.3 (iii), we obtain that the generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_2}(-1,-1)$ if and only if $\overline{l_{2n+2}} = \overline{0}$ (in $\mathbb{Z}_2$) if and only if $n \equiv 2 \pmod 3$.

Proposition 3.3. *Let p, q be two integers and let n be an arbitrary positive integer. Then, all generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_3}(-1,-1)$.*

Proof. Applying Proposition 3.1 we obtain that $n\left(G_n^{p,q}\right) = \overline{0}$ in $\mathbb{Z}_3$, $(\forall)\, n \in \mathbb{N}$. This implies that all generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_3}(-1,-1)$.

Proposition 3.4. *Let p, q be two integers and let n be an arbitrary positive integer. Then, the following statements are true:*

i) *if* $p\equiv 0\ (mod\ 5)$, *then all generalized Fibonacci-Lucas quaternions* $G_n^{p,q}$ *are zero divisors in the quaternion ring* $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$;
ii) *if* 5 *does not divide* p and $p\equiv q\ (mod\ 5)$, *then the generalized Fibonacci-Lucas quaternions* $G_n^{p,q}$ *are zero divisors in the quaternion ring* $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$ *if and only if* $n\equiv 4\ (mod\ 5)$;
iii) *if* 5 *does not divide* p and $p+q\equiv 0\ (mod\ 5)$, *then the generalized Fibonacci-Lucas quaternions* $G_n^{p,q}$ *are zero divisors in the quaternion ring* $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$ *if and only if* $n\equiv 0\ (mod\ 5)$;
iv) *if* $p\equiv 1\ (mod\ 5)$, $q\equiv 2\ (mod\ 5)$ *or if* $p\equiv 3\ (mod\ 5)$, $q\equiv 1\ (mod\ 5)$ *or if* $p\equiv 2\ (mod\ 5)$, $q\equiv 4\ (mod\ 5)$ *or if* $p\equiv 4\ (mod\ 5)$, $q\equiv 3\ (mod\ 5)$,*then the generalized Fibonacci-Lucas quaternions* $G_n^{p,q}$ *are zero divisors in the quaternion ring* $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$ *if and only if* $n\equiv 1\ (mod\ 5)$;
v) *if* $p\equiv 2\ (mod\ 5)$, $q\equiv 1\ (mod\ 5)$ *or if* $p\equiv 3\ (mod\ 5)$, $q\equiv 4\ (mod\ 5)$ *or if* $p\equiv 4\ (mod\ 5)$, $q\equiv 2\ (mod\ 5)$ *or if* $p\equiv 1\ (mod\ 5)$, $q\equiv 3\ (mod\ 5)$, *then the generalized Fibonacci-Lucas quaternions* $G_n^{p,q}$ *are zero divisors in the quaternion ring* $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$ *if and only if* $n\equiv 3\ (mod\ 5)$.
vi) *if* 5 *does not divide* p *and* $q\equiv 0\ (mod\ 5)$, *then the generalized Fibonacci-Lucas quaternions* $G_n^{p,q}$ *are zero divisors in the quaternion ring* $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$ *if and only if* $n\equiv 2\ (mod\ 5)$.

Proof. i) If $p\equiv 0\ (\text{mod } 5)$ according to Proposition 3.1 we obtain that $n\left(G_n^{p,q}\right)=\overline{0}$ in $\mathbb{Z}_5$, $(\forall)\ n\in\mathbb{N}$. This implies that all generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$.
ii) Applying Proposition 3.1, we have: $n\left(G_n^{p,q}\right)=\overline{3p^2f_{2n+1}+6pq\left(f_{2n+1}+f_{2n+3}\right)}$ in $\mathbb{Z}_5$. Using the recurrence of Fibonacci sequence, it results that:

$$n\left(G_n^{p,q}\right)=\overline{p\cdot\left[\left(3p+2q\right)f_{2n+1}+qf_{2n+2}\right]}\ in\ \mathbb{Z}_5.$$

The generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$ if and only if $n\left(G_n^{p,q}\right)=\overline{0}$ (in $\mathbb{Z}_5$). Since 5 does not divide p, the $n\left(G_n^{p,q}\right)=\overline{0}$ (in $\mathbb{Z}_5$) if and only if $(3p+2q)f_{2n+1}+qf_{2n+2}\equiv 0\ (\text{mod } 5)$. Since $p\equiv q\ (\text{mod } 5)$, we obtain that the congruence $(3p+2q)f_{2n+1}+qf_{2n+2}\equiv 0\ (\text{mod } 5)$ is equivalent with $qf_{2n+2}\equiv 0\ (\text{mod } 5)$ if and only in $f_{2n+2}\equiv 0\ (\text{mod } 5)$. According to Proposition 1.5, the last congruence is equivalent with $2n+2\equiv 0\ (\text{mod } 5)$ if and only if $n\equiv 4\ (\text{mod } 5)$.
iii) If 5 does not divide p, similarly as in ii) the generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$ if and only if $(3p+2q)f_{2n+1}+qf_{2n+2}\equiv 0\ (\text{mod } 5)$. Using the fact that $p+q\equiv 0\ (\text{mod } 5)$ and the recurrence of Fibonacci sequence, the last congruence is equivalent with $qf_{2n}\equiv 0\ (\text{mod } 5)$ if and only if $f_{2n}\equiv 0\ (\text{mod } 5)$ if and only if $n\equiv 0\ (\text{mod } 5)$ (according to Proposition 1.5).
iv) Similarly as in ii), the generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$ if and only if $(3p+2q)f_{2n+1}+qf_{2n+2}\equiv 0\ (\text{mod } 5)$. We denote $E=(3p+2q)f_{2n+1}+qf_{2n+2}$. We have:

$$E \equiv \begin{cases} 2f_{2n+1} + 2f_{2n+2} \ (mod\ 5),\ if\ p \equiv 1\ (mod\ 5),\ q \equiv 2\ (mod\ 5) \\ f_{2n+1} + f_{2n+2} \ (mod\ 5),\ if\ p \equiv 3\ (mod\ 5),\ q \equiv 1\ (mod\ 5) \\ 4f_{2n+1} + 4f_{2n+2} \ (mod\ 5),\ if\ p \equiv 2\ (mod\ 5),\ q \equiv 4\ (mod\ 5) \\ 3f_{2n+1} + 3f_{2n+2} \ (mod\ 5),\ if\ p \equiv 4\ (mod\ 5),\ q \equiv 3\ (mod\ 5). \end{cases}$$

This is equivalent with:

$$E \equiv \begin{cases} 2f_{2n+3} \ (mod\ 5),\ if\ p \equiv 1\ (mod\ 5),\ q \equiv 2\ (mod\ 5) \\ f_{2n+3} \ (mod\ 5),\ if\ p \equiv 3\ (mod\ 5),\ q \equiv 1\ (mod\ 5) \\ 4f_{2n+3} \ (mod\ 5),\ if\ p \equiv 2\ (mod\ 5),\ q \equiv 4\ (mod\ 5) \\ 3f_{2n+3} \ (mod\ 5),\ if\ p \equiv 4\ (mod\ 5),\ q \equiv 3\ (mod\ 5). \end{cases}$$

For to have $E \equiv 0 \pmod 5$ it is necessary and sufficient to have $f_{2n+3} \equiv 0 \pmod 5$. According to Proposition 1.5, the last congruence is equivalent with $2n + 3 \equiv 0 \pmod 5$ if and only if $n \equiv 1 \pmod 5$.
v) Similarly as in iv), we have:

$$E \equiv \begin{cases} 3f_{2n+1} + f_{2n+2} \ (mod\ 5),\ if\ p \equiv 2\ (mod\ 5),\ q \equiv 1\ (mod\ 5) \\ 2f_{2n+1} + 4f_{2n+2} \ (mod\ 5),\ if\ p \equiv 3\ (mod\ 5),\ q \equiv 4\ (mod\ 5) \\ f_{2n+1} + 2f_{2n+2} \ (mod\ 5),\ if\ p \equiv 4\ (mod\ 5),\ q \equiv 2\ (mod\ 5) \\ 4f_{2n+1} + 3f_{2n+2} \ (mod\ 5),\ if\ p \equiv 1\ (mod\ 5),\ q \equiv 3\ (mod\ 5). \end{cases}$$

Using Fibonacci recurrence, we obtain:

$$E \equiv \begin{cases} f_{2n} + 4f_{2n+1} \ (mod\ 5),\ if\ p \equiv 2\ (mod\ 5),\ q \equiv 1\ (mod\ 5) \\ 2f_{2n+3} + 2f_{2n+2} \ (mod\ 5),\ if\ p \equiv 3\ (mod\ 5),\ q \equiv 4\ (mod\ 5) \\ f_{2n+3} + f_{2n+2} \ (mod\ 5),\ if\ p \equiv 4\ (mod\ 5),\ q \equiv 2\ (mod\ 5) \\ -f_{2n+1} - 2f_{2n+2} \ (mod\ 5),\ if\ p \equiv 1\ (mod\ 5),\ q \equiv 3\ (mod\ 5). \end{cases}$$

This is equivalent with:

$$E \equiv \begin{cases} 4f_{2n-1} \ (mod\ 5),\ if\ p \equiv 2\ (mod\ 5),\ q \equiv 1\ (mod\ 5) \\ 2f_{2n+4} \ (mod\ 5),\ if\ p \equiv 3\ (mod\ 5),\ q \equiv 4\ (mod\ 5) \\ f_{2n+4} \ (mod\ 5),\ if\ p \equiv 4\ (mod\ 5),\ q \equiv 2\ (mod\ 5) \\ -f_{2n+4} \ (mod\ 5),\ if\ p \equiv 1\ (mod\ 5),\ q \equiv 3\ (mod\ 5). \end{cases}$$

Similarly as in ii), the generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ are zero divisors in the quaternion ring $\mathbb{H}_{\mathbb{Z}_5}(-1,-1)$ if and only if $E \equiv 0 \pmod 5$. This is equivalent with $f_{2n-1} \equiv 0 \pmod 5$, respectively $f_{2n+4} \equiv 0 \pmod 5$. According to Proposition 1.5, each of these two congruences is equivalent to $n \equiv 3 \pmod 5$.
vi) We get $E \equiv 3pf_{2n+1} \pmod 5$. For to have $E \equiv 0 \pmod 5$ it is necessary and sufficient to have $f_{2n+1} \equiv 0 \pmod 5$. According to Proposition 1.5, the last congruence is equivalent with $2n + 1 \equiv 0 \pmod 5$ if and only if $n \equiv 2 \pmod 5$.

Now, we consider $r = p \neq q$, $p > 5$ and we obtain the following result:

Theorem 3.1. *Let p, q be two prime positive integers, $p \neq q$, $p > 5$ and let n*

be an arbitrary positive integer. Then, the generalized Fibonacci-Lucas quaternions $G_n^{p,q}$ *are zero divisors in the quaternion ring* $\mathbb{H}_{\mathbb{Z}_p}(-1,-1)$ *if and only if* $n = \frac{(2l+1)\cdot z(p)-1}{2} - 1$, *where* $l \in \mathbb{N}$.

Proof. Applying Proposition 3.1 and the fact that $r = p$, it results that $\mathbf{n}(G_n^{p,q}) = \overline{15q^2 f_{2n+3}}$ in $\mathbb{Z}_p$. Since p, q are primes, $p \neq q$, $p > 5$, we obtain that $\mathbf{n}(G_n^{p,q}) = \overline{0}$ (in $\mathbb{Z}_p$) if and only if $p | f_{2n+3}$. Similarly with the proof of Theorem 2.2 we obtain that $p | f_{2n+3}$ is equivalent with $n = \frac{(2l+1)\cdot z(p)-1}{2} - 1$, where $l \in \mathbb{N}$.

Now we consider $r = q \neq p$, $q > 5$ and we obtain the following result:

Theorem 3.2. *Let* p, q *be two prime positive integers,* $p \neq q$, $q > 5$ *and let* n *be an arbitrary positive integer. Then, the generalized Fibonacci-Lucas quaternions* $G_n^{p,q}$ *are zero divisors in the quaternion ring* $\mathbb{H}_{\mathbb{Z}_q}(-1,-1)$ *if and only if* $n = \frac{(2l+1)\cdot z(q)-1}{2}$, *where* $l \in \mathbb{N}$.

Proof. Applying Proposition 3.1 and the fact that $r = q$, it results that $\mathbf{n}(G_n^{p,q}) = \overline{3p^2 f_{2n+1}}$ in $\mathbb{Z}_q$. Since p, q are primes, $p \neq q$, $q > 5$, we obtain that $\mathbf{n}(G_n^{p,q}) = \overline{0}$ (in $\mathbb{Z}_q$) if and only if $q | f_{2n+1}$. Similarly with the proof of Theorem 2.2 we obtain that $q | f_{2n+1}$ is equivalent with $n = \frac{(2l+1)\cdot z(q)-1}{2}$, where $l \in \mathbb{N}$.

References

[Ak, Ko, To; 14] M. Akyigit, HH Kosal, M. Tosun, *Fibonacci Generalized Quaternions* , Adv. Appl. Clifford Algebras, vol. **24**, issue: 3 (2014), p. 631-641

[Da, Dr; 70] D. E. Daykin, L. A. G. Dresel, *Factorization of Fibonacci numbers*, The Fibonacci Quarterly 8, No. 1 (1970).

[Fl, Sa; 15] C. Flaut, D. Savin, *Quaternion algebras and generalized Fibonacci-Lucas quaternions*, Adv. Appl. Clifford Algebras, vol. **25** issue 4 (2015), p. 853-862.

[Gr, Mi, Ma; 15] J. M. Grau, C. Miguel and A. M. Oller-Marcen, *On the Structure of Quaternion Rings Over* $\mathbb{Z}/n\mathbb{Z}$, Adv. Appl. Clifford Algebras, vol. **25**, issue: 4 (2015), p. 875-887

[Ho; 61] A. F. Horadam, *A Generalized Fibonacci Sequence*, Amer. Math. Monthly, **68**(1961), p. 455-459.

[Ho; 63] A. F. Horadam, *Complex Fibonacci Numbers and Fibonacci Quaternions*, Amer. Math. Monthly, **70**(1963), 289-291.

[Lam; 04] T. Y. Lam, *Introduction to Quadratic Forms over Fields,* American Mathematical Society, 2004.

[Mi, Se; 11] C. J. Miguel, R. Serodio, *On the structure of quaternion rings over* $\mathbb{Z}_p$, International Journal of Algebra, vol.5, 2011, no.27, 1313-1325.
[Po, Ki, Ke; 16] E. Polatli, C. Kizilates, S. Kesim, *On Split k-Fibonacci and k-Lucas Quaternions*, Adv. Appl. Clifford Algebras, vol. **26** issue 1 (2016), p. 353-362
[Su] Sun, Z. H. Congruences for Fibonacci numbers, http://www.hytc.cn/xsjl/szh
[Su; 92] Sun, Z. H., Z.W. Sun. Fibonacci numbers and Fermats last theorem. Acta Arithemtica, Vol. 60, 1992, 371388.
[Ta; Ya; 15] D. Tasci, F. Yalcin, *Fibonacci-p Quaternions*, Adv. Appl. Clifford Algebras, vol. **25** issue 1 (2015), p. 245-254.
[Fib.] http://www.maths.surrey.ac.uk/hosted-sites/R.Knott/Fibonacci/fib.html
[Yü, Ay; 16] S. Yüce, F.T. Aydin, *A New Aspect of Dual Fibonacci Quaternions,* accepted in Adv. Appl. Clifford Algebras (http://link.springer.com/article/10.1007/s00006-015-0619-9).

Diana SAVIN
Faculty of Mathematics and Computer Science,
Ovidius University,
Bd. Mamaia 124, 900527, CONSTANTA, ROMANIA
http://www.univ-ovidius.ro/math/
e-mail: savin.diana@univ-ovidius.ro, dianet72@yahoo.com